INDEX TO VOLUME XVIII

New species in blackface type

Abacá, root knot of, 861–867

Abies, amabilis, host of needle rusts, 705–708; arizonica, host of Peridermium pustulatum, 705–708; balsamea, host of needle rusts, 705–708; concolor, host of needle rusts, 705–708; fraseri, host of needle rusts, 705–708; grandis, host of needle rusts, 705–708; lasiocarpa, host of needle rusts, 705–708; magnifica, host of needle rusts, 705–708; nobilis, host of needle rusts, 705–708; species, conspectus of needle rusts on, 705–708; venusta, host of Peridermium pseudobalsameum, 705–708

Abstracts, of papers presented at the nineteenth annual meeting of the American Phytopathological Society, Nashville, Tenn., December 28, 1927, to January 1, 1928, 127–156; of papers read at the ninth annual meeting, Canadian Division of the American Phytopathological Society, 477–481

Acanthorhynchus vaccinii, causing cranberry storage rot, 809–814

Accuracy in comparing concentrations of tobacco-mosaic virus, 132

Acer, pennsylvanicum, lesions produced by Septoria acerina, 734; rubrum, lesions produced by Rhytisma acerinum, 744

Actinomycete, causing soil rot or pox in sweet potatoes, 152

ADAMS, J. F., An Actinomycete the cause of soil rot or pox in sweet potatoes, 152

Aecia, in simple (monosporidial) pustules, 1008–1010; occurrence of, in certain rust fungi, 1005–1015; occurrence under natural conditions of pycnia, unaccompanied by, 1010–1014; of white-pine blister rust, permanent mounts of, 1029

Aecial host, in relation to severity of stem-rust epidemics, 105–108

Aecidium, onobrichydis, 665; polygalinum, 665

Agar, carrot, for Pythium, 710

Agglutinins, production of, by phytopathogenic bacteria, 277–288

Alfalfa, white spot of, caused by irrigation, 136–137

Alkalinity, soil, clubroot in relation to, 301–306

Alternaria, citri, 949; solani, lesions produced on Lycopersicum esculentum, 740; solani, lesions produced on Solanum tuberosum, 739–740; sp., relation to apple rots, 565

American Phytopathological Society, abstracts of papers presented at Nashville meeting, Dec. 28, 1927, to Jan. 1, 1928, 127–156; Canadian Division, abstracts of papers read at, 477–481; report of nineteenth annual meeting, 463–474; report of ninth annual meeting Canadian Division, 477–481

Ampelopsis tricuspidata, lesions produced by Guignardia bidwellii, 746–747

ANDERSON, H. W., Bacteriophage of Bacterium pruni, 144

ANDERSON, P. J., Soil reaction and black root rot of tobacco, 131

Andropogon, rust on, 665

Angular leaf spot, 951

Antheridia, paragynous, of Phytophthora spp., 149–150

Anthracnose of snowberry, 797–801

Antirrhinum majus, Puccinia antirrhini on, experimental control of teliospore and urediniospore formation, 176–178, 183–184, 192–194

Antiserum, production of, against phytopathogenic bacteria, 278–288

Aphanomyces, cochlioides, n. sp., on sugar beets in the United States, 149; euteiches, pathogenicity for conifers, 366

Aphis, transmission of cucumber mosaic, 143

Apium graveolens, lesions produced by Septoria petroselini, 731

Aplanobacter stizolobii, see Bacterium stizolobii, 460

Apparatus, constant-temperature and humidity chambers, 227–238; for isolating single spores, 1017–1021

Apple, and quince trees, nursery, overgrowths and hairy root on, 127; blotch, cycle of infection in, 145; correlation of the wound overgrowth and crown gall of, 128; grafting as a further means of preventing callus knots on, 127–128; grafts, history of development of wound overgrowths on, and of the influence of wrappers on their suppression, 128; root and crown injury, to trees, 547–551; rots, caused by some fungi of the Stemphylium type, 565–578; scab, control with certain sulphur fungicides, 146–147; scab of, fall applications of fungicides in relation to control of, 146; scab, see Venturia inaequalis; sporotrichum fruit spot of, 145; studies of black root of, 145

Apples, copper sulphate as remedy for exanthema in, 449–454

Apricot, scab caused by Cladosporium, 314–315

Armillaria mellea in mines and wells, 948

ARTHUR, J. C., Progress of rust studies, 659–674

Asci, apple scab, effect of environment upon, 375–418

Ascomycetous leaf spot of cowpea, 701–704

Asparagus officinale, Puccinia asparagi on, experimental control of teliospore and urediniospore formation, 180, 181–183

Aspergilli, investigations of, by serological methods, 148; serological studies of, 691–696

Aspergillus, attacking mealybugs in insectaries in southern California, 948; niger, 949; niger, reaction of plant stems to the products of, 604–605

Aster, China, Fusarium wilt of, 150; yellows, studies on host range of, 156

Asterocystis racidis, 949–950

Athyrium cyclosorum, host of Uredinopsis copelandi, 705–708

Attenuation of plant viruses, 156

Bacillus, aroideae, humidity relations of, 827–829; aroideae, inoculation into various hosts, 832–836; aroideae, pathogenicity to potato tubers, 819–838; aroideae, temperature relations of, 822–827; carotovorus, humidity relations of, 827–829; carotovorus, inoculation into various hosts, 832–836; carotovorus, pathogenicity to potato tubers, 819–838; carotovorus, temperature relations of, 822–827; mesentericus, humidity relation of, 827–829; mesentericus, inoculation into various hosts, 832–836; mesentericus, pathogenicity to potato tubers, 819–838; mesentericus, temperature relations of, 822–827; phytophthorus, humidity relations of, 827–829; phytophthorus, inoculation into various hosts, 832–836; phytophthorus, pathogenicity to potato tubers, 819–838; phytophthorus, temperature relations of, 822–827; tracheiphilus, methyl violet for staining, 803–807

Bacteria, demonstration in plant tissues by means of Giemsa stain, 803–807; phytopathogenic, production of agglutinins by, 277–288

Bacterial, disease of broad beans, 136; redstripe disease of sugar cane in countries of the Pacific, 945

Bacteriophage of Bacterium pruni, 144

Bacteriosis of Oleander in California, 503–518

Bacterium, cerasus, 952; citriputeale, 952; **maculicola,** (McC.) nom. emend. syn. Bact. maculicolum, 460; maculicolum, production of antisera against, 278–288; maculicolum, see Bact. maculicola; pruni, bacteriophage of, 144; **stizolobii** (Wolf), comb. nov. syn. Aplanobacter stizolobii, 460; syringae, 952; tumefaciens, biology, 332–339; tumefaciens, distribution in nature, 339–342; tumefaciens, method and rate of migration in tomato, 129; tumefaciens, migration in tissue of tomato plants, 553–564; tumefaciens, see Pseudomonas tumefaciens

BAILEY, D. L., and F. J. GREANEY, see GREANEY, F. J.; Sulphur dusting for the control of leaf and stem rust in Manitoba: Field trials with horse-drawn and aeroplane dusters, 480

BAIN HENRY F., see STEVENS, NEIL E.

Balsam firs, conspectus of needle rusts on, 705–708
Banana, Chinese, "finger-tip" disease, 531–538
BANFIELD, W. M., see RIKER, A. J.; Studies on the life history of the crown-gall organism, 128–129
Barley, control of stripe disease, 477–478; dwarf leaf rust of, in western Canada, 481; hullness, control of covered smut and stripe on, 139; plant, powdery mildew, 901–910; stripe, experiments on control of, 295–300
BARRETT, J. T., Phytophthora in relation to crown rot of walnut, 948–949
BARRUS, M. F., and JAMES G. HORSFALL, Preliminary note on snowberry anthracnose, 797–801
BARSS, H. P., A question of correct usage, 460
Bean, mosaic of, transmission, 155; susceptibility to Bacillus carotovorus, B. phytophthorus, and B. aroideae, 834–835; Uromyces appendiculatus on, experimental control of teliospore and uredinio-spore formation, 163–169, 183, 184–187, 206
Beans, broad, a bacterial disease of, 136
Beech, mottle-leaf disease of, 151–152
BEECHER, F. S., Measurements of total daily sunlight intensity with reference to the ecology of plant diseases, 951; see SHAPOVALOV, M.
Beet, curly top, on celery and spinach, 709–710; sugar, predisposition of, to late root rot, 263–276; sugar, transmission of curly-top virus through different solutions, 675–679
Beets, sugar, filtration experiments on curly top of, 681–690
BENSAUDE, M., and G. W. KEITT, Comparative studies of certain Cladosporium diseases of stone fruits, 313–329
Beta, vulgaris, 939; lesions produced by Cercospora beticola, 728–730
BIJL, P. A. van der, Plantsiektes hul oorsaak en bestryding, review of, 1023
Biography, of BURGER, OWEN F., 627–630
Biology, life history of Erostrotheca multiformis, perfect stage of Cladosporium album, 839–846; life history of Sclerotium gladioli, 526–527; of Bacterium tumefaciens, 332–339
Biometrical studies on the variation of physiologic forms of Puccinia graminis tritici and the effects of ecological factors on the susceptibility of wheat varieties, 7–123
BISHOP, C. J., see DAVIS, A. R.
Black leg, potato, transmisison by seed-corn maggot, 459
Back scab, or warty disease, of potato, in Belgium, 126
"Black tip," disease of Chinese banana, 531–538
Blast, citrus, 952; in oats, 948
Blastophaga psenes, 931–932, 938
Blight, brown, disease of lettuce, 949; fire, disinfectants in eradication work, 710–711; Pahala, of sugar cane, effect of manganese deficiency on, 775–786; Phytophthora of peony, 881–900; seedling, of oats, 480; thread, further observations on, 147–148; tip caused by Phytophthora, 882–883
Blight-resistant potatoes, 483–502
Blister rust of white pine, see Cronartium ribicola
BLOOD, H. L., A "streak" of tomatoes produced by a disturbing principle from apparently healthy potatoes in combination with tomato mosaic virus, 311
Blotch, of apple, cycle of infection in, 145
BONDE, REINER, The transmission of potato black leg by the seed-corn maggot in Maine, 459
Book review, Die Krankheiten der Obstbäume und Obststräucher, by E. EWERT, 715; A list of Japanese fungi hitherto unknown, 1023; Plantsiektes hul oorsaak en bestryding, 1023
Bordeaux mixture, preventing stem rot of sweet potatoes, 152–153
Botrytis, 881
BOYCE, A. M., and H. S. FAWCETT, An Aspergillus attacking mealybugs in insectaries in southern California, 948
BOYCE, J. S., A conspectus of needle rusts on balsam firs in North America, 705–708; A possible alternate stage of Puc-

ciniastrum myrtilli (Schum.) Arth., 623–625

BOYD, O. C., Comparison of various disinfectants in the treatment of sweet potatoes for black-rot control, a progress report, 153–154; Preliminary report on a new leaf spot of pecan, 133–134

BREWBAKER, H. E., see STAKMAN, E. C.

BRIERLEY, PHILIP, Pathogenicity of Bacillus mesentericus, B. aroideae, B. carotovorus, and B. phytophthorus to potato tubers, 819–838

BROWN, A. M., and M. NEWTON, The dwarf leaf rust of barley in western Canada (Puccinia anomala Rostr.), 481; see NEWTON, M.

BROWN, H. D., see STAIR, E. C.

BROWN, J. G., The influence of sodium chloride in alkali soils on the susceptibility of Pima cotton to angular leaf spot caused by Phytomonas malvacera, 951

Brown rot, 931

Brown stripe, of sugar cane, 761–772; of sugar cane, caused by **Helminthosporium stenospilum**, n. sp., 135–136; of sugar cane, in Cuba, 135

BRYAN, MARY K., Verticillium wilt of heliotrope, 246

Bulbs, of ornamental plants, condition of, offered for importation into Canada, 150–151

Bunchy top, of abacá, relation of Heterodera radicicola to, 865–866

Bunt, effect of washing seed on infection, 711–713; infection, influence of soil temperature and moisture, 924–928; of wheat, control, in Nebraska, 921–929

BURGER, OWEN F., biography of, 627–630

Cabbage, Chinese, parasitized by a powdery mildew, 611–615; Fusarium-resistant, Jersey Wakefield type, 142; Iacope, improvement of, by selecting and selling, 142; susceptibility to Bacillus carotovorus, B. phytophthorus, and B. aroideae, 833–835

Caeoma americana, on Abies species, 705–708

CALINISAN, MELANIO R., see OCFEMIA, GERARDO OFFIMARIA

Calyptospora, columnaris, on Vaccinium species, 705–708; sp., on V. macrophyllum, 705–708

Canadian Division, American Phytopathological Society, report of ninth annual meeting of, 477–481

Cane, sugar, observations on mycorrhizae and their relation to root diseases of, 249–261; sugar, see sugar cane.

Capsicum sp., mosaic of, in West Africa, 155

Carrot, agar, for Pythium, 710; susceptibility to Bacillus carotovorus, B. aroideae, B. phytophthorus, 832–835

CARSNER EUBANKS, and C. F. LACKEY, Further studies on attenuation of the virus of sugar beet curly top, 951

CARTER, WALTER, An improvement in the technique for feeding homopterous insects, 246–247; Transmission of the virus of curly top of sugar beets through different solutions, 675–679

CATION, DONALD, see HOWE, MARY F.

Cauliflower, spot disease of, 460

Caulophyllum thalictroides, lesions produced by Cercospora caulophylli, 735

Celery, host of beet curly top, 709–710

Cerastium species, alternate host of a fir needle rust, 705–708

Cercospora, beticola, lesions produced on Beta vulgaria, 728–730; caulophylli, lesions produced on Caulophyllum thalictroides, 735; circumcissa, lesions produced on Prunus serotina, 737–738; menispermi, lesions produced on Menispermum canadense, 736–737; musarum, cause of "finger-tip" or "black tip" of Chinese banana, 533–536; symplocarpi, lesions produced on Symplocarpus foetidus, 736

Cereal, diseases, field method of insuring positive attack by, 617–621; smuts, physiologic specialization in, 955–1003

Cereals, diseases of, in Alberta, 481

Ceuthospora lunata, causing cranberry storage rot, 809–814

Chaleas exotica, 952

Changes, histologic, induced in leaves by leaf-spotting fungi, 717–751

CHARLES, V. K., see MARTIN, G. HAMILTON

CHEN, H. K., see PORTER, R. H.

Chenopodium murale, 951
Cherry, scab caused by Cladosporium, 315–316
Chinese cabbage, powdery mildew parasitizing, 611–615
Chlorophyll content, tobacco, effects of mosaic upon, 697–700
CHRISTENSEN, J. J., see IMMER, F. R.; see STAKMAN, E. C.
Chromosome, number, and reaction to rust and mildew in Linum, 478
CHUPP, CHAS., Clubroot in relation to soil alkalinity, 301–306
CHURCH, MARGARET B., and MARIO SCANDIFFIO, Phytophthora, Pythium, and Pythiacystis species as stock cultures, 247–248
Cicatrice, diseases in which it is definitely formed about margin of lesion, 723–731; diseases in which it is not definitely formed about edge of lesions, 731–747
Ciccinnobolus cesatii, parasite of Sphaerotheca humuli on raspberry, 789
CIFERRI, R., Preliminary observations on sugar cane mycorrhizae and their relation to root diseases, 249–261
Cirsium arvense, lesions produced by Septoria cirsii, 733; Puccinia suaveolens on, experimental control of teliospore and urediniospore formation, 169–172, 191–192, 206
Citrus, aurantium, 953; blast and some allied organisms, 952; leaves, histology of lesions due to Sphaceloma fawcetii Jenkins, 539–545; limonia, 953
Cladosporium, album, life history of, 839–844; carpophilum, 328, 946; cerasi, see Venturia cerasi; cultural morphology, 324–327; diseases, 946; diseases of stone fruits, 313–329; of tomato, control by means of forced ventilation in hotbeds, 1027–1029; strains affecting stone fruits, 327–328; temperature relations, 322–324
Climatic conditions, effect on mycorrhizae, 258–259; in relation to severity of stem-rust epidemics, 91–105
Clover, Uromyces trifolii on, experimental control of teliospore and urediniospore formation, 172–175, 188–191, 206
Clubroot, of crucifers, reaction and treatment of soils infested with Plasmodiophora brassicae, 141–142; in relation to soil alkalinity, 301–306
Coccomyces, hiemalis, lesions produced on Prunus avium, 727; hiemalis, lesions produced on Prunus cerasus, 726–727; lutescens, lesions produced on Prunus virginiana, 725–726; prunophorae, lesions produced on Prunus domestica, 723–725
Collection, photographic, of ERWIN F. SMITH, 312
Comandra, rust on, 665
Conifers, seedling damping off, 361–367
Conspectus of needle rusts on balsam firs in North America, 705–708
Control, bunt of wheat, in Nebraska, 921–929; of apple scab in relation to fall applications of fungicides, 146; of apple scab, with certain sulphur fungicides, 146–147; of barley stripe, 477–478; of barley stripe, experiments, on, 295–300; of "black tip" of Chinese banana, 536–537; of covered smut and stripe of barley by seed disinfectants, 139; of cucumber powdery mildew in greenhouses, 847–860; of dry rot of Gladiolus corms, 527–528; of tomato Cladosporium and Septoria in hotbeds, by means of forced ventilation, 1027–1029; of true tobacco mosaic, 132; of turf diseases, with mercury, 137
COOK, MELVILLE T., The gummosis of sugar cane, 135
COOPER, D. C., and C. L. PORTER, Phytophthora blight of peony, 881–900
COOPER, DELMER, Paragynous antheridia of Phytophthora spp., 149–150
Copper, fungicides, classification of, 455–458; sulphate, remedy for exanthema in prunes, apples, pears, and olives, 449–454
Coprosma baueri, 952
Corm, of Gladioli, Penicillium rot of, 151; rot of Gladiolus caused by a Penicillium, 151
Corms of ornamental plants, condition of, offered for importation into Canada, 150–151
Corn, influence of environmental factors on prevalence of smut in, 589; losses due to

smut, 599–602; method of measuring efficiency of seed disinfectants on, 140; Penicillium injury to seedlings, 239–242; Puccinia sorghi on, experimental control of teliospore and urediniospore formation, 179–180, 194–195, 206; rust of, Puccinia sorghi, 345–354

Correlation of the wound overgrowth and crown gall of apple in parts of Europe and of the United States, 128

Corticium koleroga, further observations on, 147–148

Cotton, outstanding wilt-resistant varieties of, 134; seed treatment by the dusting method, 134; wilt fungus, consideration of the pathogenicity of, 419–438

Cowpea, ascomycetous leaf spot of, 701–704

CRAIGIE, J. H., Heterothallism in the rust fungi, 147; On the occurrence of pycnia and aecia in certain rust fungi, 1005–1015; Sexual behaviour of Puccinia graminis, 479

Cranberries, storage rots, in 1927, 809–814

Cronartium, cerebrum, 671; ribicola, permanent mounts of aecia of, 1029

Cropping systems of fertilizers, influence of, on black and brown root rot of tobacco, 131

Crown, and root injury to apple trees, 547–551; elongation disease of peony, 243–244; rot of walnut, 948–949

Crown gall, of apple, correlation of, and wound overgrowth, 128; organism, studies on the life history of, 128–129; situation in England, France, and Holland, notes on, 289–294; transplants, studies on, 130

Cucumber, Chinese, resistant to mosaic, 143; control of powdery mildew on, 847–860; fruit rot and angular leaf spot, 133; mosaic, transmission of, by aphis, 143; susceptibility to Bacillus aroideae, 834–835

Cucumis melo, lesions produced by Macrosporium cucumerinum, 740–741

Cultural, characteristics, Pseudomonas savastanoi var. nerii, 513–515; characters of Sclerotium gladioli, 524–526; conditions, effect of, on stability of physiologic forms of Puccinia graminis tritici, 108–113; differences, between physiologic forms of cereal smuts, 958–972; studies of Cladosporium strains, 325–327

Cultures, stock, of Phytophthora, Pythium, and Pythiacystis, 247–248

CUNNINGHAM, H. S., A study of histologic changes induced in leaves by certain leaf-spotting fungi, 717–751; Histology of the lesions produced by Sphaceloma fawcettii Jenkins on leaves of Citrus, 539–545

Curly top, of sugar beet, 951; of sugar beet, transmission of virus through different solutions, 675–679; of sugar beets, filtration experiments on, 681–690; tomato, 709–710

Cuscuta gronovii, haustorium of, 478

Cycle of infection in apple blotch, 145

Cylindrocladium macrosporum, n. sp., cause of leaf spot of Washingtonia palm, 219–225

Cytological, phenomena in disease-resistant plants, 144; studies of plant tissues affected with mosaic diseases, 154

DAHL, A. S., see MONTEITH, JOHN, JR.

Damping off, of coniferous seedlings, 361–367

Dandelion, Puccinia taraxaci on, experimental control of teliospore and urediniospore formation, 175–176, 196–197

DAVIS, A. R., R. H. MARLOTH, and C. J. BISHOP, The inorganic nutrition of the fungi. I. The relation of calcium and boron to growth and spore formation, 949

DAVIS, W. H., A powdery mildew parasitizing Chinese cabbage, 611–615; Two physiological forms of Ustilago striaeformis (Westd.) Niessl., 149

Development of tomato yellows under different light conditions, 950

Dieback, see exanthema

DIETZ, S. M., and L. D. LEACH, The effective methods of eradicating Rhamnus species susceptible to Puccinia coronata Corda, 138

Diplodia natalensis, 949

Disease, bacterial, of broad beans, 136; crown elongation of peony, 243–244; mottle-leaf, of beech, 151–152; of potato, new and destructive, in Utah, and its relation to the potato psylla, 140–141; spike, of Santalum album and Vinca rosea, occurrence of intracellular bodies in, 815–817; warty, or black scab, of potato, in Belgium, 126

Diseases, cereal, field method of insuring positive attack, 617–621; Cladosporium, of stone fruits, 313–329; mosaic, cytological studies of plant tissues affected with, 154; of cereals, in Alberta, 481; of turf, controlled with mercury, 137; virus, of tobacco in Kentucky, 132–133

Disinfectants, controlling sweet-potato black rot, 153–154; effect of, upon germination of seeds in storage for indefinite periods after treatment, 136; in fire-blight-eradication work, 710–711; of seed corn, method of measuring relative efficiency, 140; seed, for control of covered smut and stripe of barley, 139

DOAK, K. D., The mycorrhizal fungus of Vaccinium, 148

DOOLITTLE, S. P., Soil transmission of tomato mosaic and streak in the greenhouse, 155; and M. N. WALKER, Aphis transmission of cucumber mosaic, 143

DORAN, W. L., Effect of timothy infusion of different ages on the growth of tobacco and on brown root rot of tobacco, 131–132

DOSDALL, LOUISE, A mosaic disease of Gladiolus, 215–217

Dothiorella, ribis, 949; sp., 949

DRAYTON, F. L., Progress report on the condition of bulbs and corms of ornamental plants offered for importation into Canada, 150–151

DRECHSLER, CHARLES, The occurrence of **Aphanomyces cochlioides**, n. sp., on sugar beets in the United States, 149; **Pythium arrhenomanes**, n. sp., a parasite causing maize root rot, 873–875; A species of Helminthosporium distinct from H. sacchari, causing brown stripe of sugar cane, 135–136

Dry rot of Gladiolus corms, 519–529

Dryopteris, marginalis, host of Milesia marginalis, 705–708; spinulosa, host of M. kriegerina, 705–708

DUFRÉNOY, J., Cytological studies of plant tissues affected with mosaic diseases, 154; Some cytological phenomena in disease-resistant plants, 144

DUNLAP, ALBERT A., Effects of mosaic upon the chlorophyll content of tobacco, 697–700

Dust treatments for oat smuts, 139

Dusting, cotton-seed, 134

Dwarf, disease of strawberry, 439–444

Ecological factors, effects of, on susceptibility of wheat varieties to stem rust, 81–108

Edaphic conditions, effect on mycorrhizae, 258–259; in relation to severity of stem-rust epidemics, 90–91

Effect, of disinfectants upon the germination of seeds kept in storage for indefinite periods after treatment, 136; of manganese deficiency on sugar-cane plant and its relationship to Pahala blight of sugar cane, 775–786; of mixed inoculations of certain citrus fruit-rotting organisms, 949; of washing the seed on infection of wheat by stinking smut, 711–713

Effective methods of eradicating Rhamnus species susceptible to Puccinia coronata Corda, 138

Effects of lime on cigar tobacco, 130–131; of mosaic upon the chlorophyll content of tobacco, 697–700

ELIASON, E. J., Comparative virulence of certain strains of Pythium in direct inoculation of conifers, 361–367

ELMER, O. H., Penicillium corm rot of Gladioli, 151

Endophytes, taxonomy of sugar cane mycorrhizal, 251–256

Endosepsis, 931; and its control in caprifigs, 931–938

England, crown gall situation in, 289–294

Environment, influence of, in apple scab, 375–418; influence of, on seasonal prevalence of corn smut, 589–598

Eocronartium muscicola, 952; fr. in a new locality and on a new host, 952
Epidemics, of stem rust, effect of ecological factors on, 81–108
Epidemiology, prevalence and distribution of physiologic forms of cereal smuts, 958–959, 997–999; prevalence and distribution of physiologic forms of Puccinia graminis tritici isolated from uniform rust nurseries in 1919–1923, 35–40; wheat stem rust, 369–374
Epilobium species, alternate hosts of fir-needle rusts, 705–708; hosts of Pucciniastrum epilobii, 705–708
Epiphytotics, see epidemiology
Eradicating species of Rhamnus, 138
Eradication, fire-blight, disinfectants in, 710–711
Eriksson prizes, announcement of, 1029-1030
Erostrotheca, multiformis, cultural studies of, 841–843; multiformis, life history of, 839–846
Errata, correction for "Virus mixtures that may not be detected in young tobacco plants," by H. H. McKinney, Phytopath. *16*: 893, 313
Erysiphaceae, 902
Erysiphe, cichoracearum, 904; cichoracearum, control of, 847–860; graminis, 901–904, 909; graminis, method of epidemic induction, 617–621; polygoni, parasitizing Chinese cabbage, 611–615
Etiology, of abacá root knot, 863–865; of decay in apples, 576; of dry rot of Gladiolus, 521–524; of Oleander bacteriosis, 512–513
Euonymus japonicus, 904
Euttetix tenellus, carrier of sugar-beet root rot, 263; transmission of sugar-beet curly top by, 675–679, 681–690
Ewert, E., Die Krankheiten der Obstbäume und Obststräucher, book review, 715
Exanthema, copper sulphate as remedy for, in prunes, apples, pears, and olives, 449–454
Experimental work with the transmission of bean mosaic, 155
Experiments, and observations on the control of true tobacco mosaic, 132; on control of barley stripe, 295–300; with tomato streak, 154
Eye-end rot, 931
Eye spot of sugar cane, caused by Helminthosporium sacchari, 135–136; H. ocellum causes, 757–761, 763–774; in Cuba, 135

Factors important in the development of perithecia of Venturia inaequalis, 145–146
Fajardo, T. G., Progress on experimental work with the transmission of bean mosaic, 155
Fall applications of fungicides in relation to apple-scab control, 146
Faris, James A., Brown stripe of sugar cane in Cuba, 135; Sugar-cane eye spot in Cuba, 135; Three Helminthosporium diseases of sugar cane, 753–774
Fawcett, H. S., see Boyce, A. M.; see Savastano, G.
Felix bulbifera, host of Uredinopsis copelandi, 705–708
Fertilizers and cropping systems, influence of, on black and brown root rot of tobacco, 131
Field meeting, Southern Section, American Phytopathological Society, notice of, 460
Filtration experiments on curly top of sugar beets, 681–690
"Finger-tip," disease of Chinese banana, 531–538
Fire blight, disinfectants in eradication work, 710–711
Firs, balsam, conspectus of needle rusts on, 705–708
Fisk, Vernon C., see Melhus, I. E.
Forced ventilation as a means of controlling tomato Cladosporium and Septoria in hotbeds, 1027–1029
Fragaria virginiana, lesions produced by Mycosphaerella fragariae, 744
France, crown-gall situation in, 289–294
Fraxinus, americana, lesions produced by Phyllosticta fraxinii, 741; floribunda, 952

FROMME, F. D., The effect of washing the seed on infection of wheat by stinking smut, 711–713; see WINGARD, S. A.; Studies of black root rot of apple, 145
Fruit rot, of cucumber, and angular leaf spot, 133
Fruit-rotting organisms, 949
Fruit spot, sporotrichum, of apple, 145
Fruits, stone, Cladosporium diseases, 313–329
Fungi, inorganic nutrition of, 949; leaf-spotting, histologic changes induced in leaves by, 717–751; rust, occurrence of pycnia and aecia in, 1005–1015
Fungicidal, activity, of sulphur, increased by oxidizing agents, 1026–1027; property of sulphur, factors affecting, 147
Fungicides, certain sulphur, in control of apple scab, 146–147; copper, classification of, 455–458; fall applications of, in relation to apple-scab control, 146; lime-sulphur, action, 355–360
Fungous products, reaction of plant stems to, 603–610
Furfural, penetration of, in plant tissues, 152
Further studies on attenuation of the virus of sugar-beet curly top, 951
Fusaria in the herbarium of Pathological Collections, B. P. I., U. S. Dept. Agr., 148–149
Fusarium, 941; batatatis, sweet-potato stem rot prevented by Bordeaux mixture, 152–153; culmorum, on oat seedlings, 480; eumartii, varietal susceptibility of potatoes to, 307–309; lycopersici, reaction of plant stems to products of, 604–605; moniliforme var. fici, 931, 936, 938; niveum, cause of watermelon wilt, 143–144; niveum, varietal resistance of watermelons to, 144; sp., 952; **tumidum,** n. sp., on Sorothamnus scoparius, 148; vasinfectum, pathogenicity of, 419–438; wilt and stem-end rot, varietal susceptibility of potatoes to, 307–309; wilt of China aster, 150
Fusarium-resistant cabbage of Jersey Wakefield type, 142
Fusicoccum, fraxini (Kabat. et Bulak), n. com., on Fraxinus excelsior, 148; putrefaciens, cause of cranberry storage rot, 809–814

GAINES, E. F., New physiologic forms of Tilletia levis and T. tritici, 579–588; New physiologic forms of T. tritici in wheat, 139
GALLOWAY, BEVERLY T., FLORA W. PATTERSON, 1847–1928, 877–879
GARDNER, MAX W., Sporotrichum fruit spot of apple, 145
Germination, effect of disinfectants upon, of seeds kept in storage, 136; of Puccinia glumarum, optimum temperature for, 481
Gibberella quinqueseptata, n. sp., on Cannabis sativa, 148
Giemsa stain, demonstration of bacteria in plant tissues by means of, 803–807
GILLIGAN, G. M., see HOLLAND, E. B.
Gladioli, Penicillium corm rot of, 151
Gladiolus corms, dry rot, 519–529; mosaic disease of, 215–217
Glomerella rufomaculans, causing cranberry storage rot, 809–814
Gnomonia leptostyla, lesions produced on Juglans cinerea, 746
GOLDSWORTHY, M. C., The fungicidal action of liquid lime sulphur, 355–360; Production of agglutinins by phytopathogenic bacteria, 277–288
GORDON, W. L., Physiologic forms of Puccinia graminis avenae Eriks. & Henn., in Canada, 479–480
GOSS, R. W., Transmission of potato spindle tuber by grasshoppers (Locustidae), 140, 445–448; Varietal susceptibility of potatoes to Fusarium wilt and stem-end rot, 307–309
GOULDEN, C. H., K. W. NEATBY, and J. N. WELSH, The inheritance of resistance to Puccinia graminis tritici in a cross between two varieties of Triticum vulgare, 631–658
Grafting, as a further means of preventing callus knots on apple, 127–128
Grafts, apple, history of development of wound overgrowths on and of the influence of wrappers on their suppression, 128

Grasshopper (Locustidae), transmitting potato spindle tuber, 140
Grasshoppers, transmission of potato spindle tuber by Locustidae, 445–448
Grated carrot agar favorable for studies of Pythium, 710
GREANEY, F. J., and D. L. BAILEY, see BAILEY, D. L.; Sulphur dusting for the control of leaf and stem rust in Manitoba: Winnipeg small-plot experiments, 480
GUBA, E. F., Control of cucumber powdery mildew in greenhouses, 847–860
Guignardia, bidwellii, lesions produced on Ampelopsis tricuspidata, 746–747; vaccinii, causing cranberry storage rot, 809–814
Gummosis of sugar cane, 135
GUTERMAN, E. F., Preliminary report on mechanical transmission of the mosaic of Lilium auratum, 1025–1026
Gymnoconia interstitialis, 671
Gymnosporangium sp., heterothallism in, 1005–1015

Hairy root, and overgrowths, on nursery-apple and quince trees, 127
HAMILTON, J. M., and G. W. KEITT, Certain sulphur fungicides in the control of apple scab, 146–147
HANNA, W. F., A simple apparatus for isolating single spores, 1017–1021
Hansen, Emil Ch., Medal, 1928, 125
HANSEN, H. N., Endosepsis and its control in caprifigs, 931–938
HARA, K., see SHIRAI, M.
Haustorium of Cuscuta gronovii, 478
Heliotrope, Verticillium wilt of, 246
Helminthosporiose of sugar cane, 754–758
Helminthosporium, diseases, three, of sugar cane, 753–774; gramineum, experiments on control of, 295–300; **ocellum**, n. sp., causing eye spot of sugar cane, 757–761, 763–772; sacchari, causing helminthosporiose of sugar cane, 754–758; sacchari, the cause of eye spot of sugar cane, 135–136; sativum, method of epidemic induction, 617–621; stenospilum, causing brown stripe of sugar cane in Cuba, 761–772; **stenospilum**, n. sp., causing brown stripe of sugar cane, 135–136; teres, method of epidemic induction, 617–621
Hemp, Manila, root knot of, 861–867
HENRY, A. W., Reaction of Linum species of various chromosome numbers to rust and powdery mildew, 478
Heterodera radicicola, description, 863–865; on abacá, 861–867; relation to bunchy top of abacá, 865–866
Heteroecism, a possible alternate stage of Pucciniastrum myrtilli, 623–625
Heterothallism, evidence of, in Puccinia graminis, 1005–1008; in certain rust fungi, 1005–1015; in the rust fungi, 147
HIENTON, T. E., see STAIR, E. C.
HILL, J. BEN, The method and rate of migration of Bacterium tumefaciens in tomato, 129; The migration of Bact. tumefaciens in the tissue of tomato plants, 553–564
Histologic changes induced in leaves by leaf-spotting fungi, 717–751
Histology of scab lesions on Citrus leaves, 539–545
Holland, crown-gall situation in, 289–294
HOLLAND, E. B., and G. B. GILLIGAN, Classification of copper fungicides, 455–458
HOLMES, FRANCIS O., Accuracy in comparing various concentrations of tobacco-mosaic virus, 132; Monochromatic light photography in the study of mosaic diseases, 154
Hordeum vulgare, 905
HORSFALL, JAMES G., see BARRUS, M. F.
Host, range, see parasitism
Hotbeds, forced ventilation as a means of controlling tomato Cladosporium and Septoria in, 1027–1029
HOTSON, J. W., Armillaria mellea in mines and wells, 948; Rusts of the Pacific Northwest, 948
HOWARD, FRANK L., Disinfectants in fire-blight-eradication work, 710–711
HOWE, MARY F., and DONALD CATION, The penetration of furfural in plant tissues, 152
HOWITT, J. E., and R. E. STONE, Results

of experiments on the control of barley stripe, 477–478
Humidity, chambers, constant-control, 227–238; effect of, on bacterial tuber rots, 827–829
HUNTER, H. A., see JEHLE, R. A.
HURSH, C. R., The reactions of plant stems to fungous products, 603–610
Hyalopsora aspidiotus, on Phegopteris dryopteris, 705–708
Hylemyia cilicrura, transmission of potato black leg by, in Maine, 459
Hypnaceae, 952

Ilex verticillata, lesions produced by Rhytisma ilices-canadensis, 743
IMMER, F. R., and J. J. CHRISTENSEN, Determination of losses due to smut infections in selfed lines of corn, 599–602; Influence of environmental factors on the seasonal prevalence of corn smut, 589–598
Infection, cycle of, in apple blotch, 145; of wheat by stinking smut, effect of washing the seed upon, 711–713
Inheritance of resistance, to blast in oats, 948; to Puccinia graminis tritici in a cross between two varieties of Triticum vulgare, 631–658; to P. sorghi in maize, 138
Inhibition of enzymatic action as a possible factor in the resistance of plants to disease, 953
Injury, leaf-hopper, to legumes, 137–138
In memoriam, 475
Inoculum, quantity of, in relation to severity of stem-rust epidemics, 81–83
Insects, carriers of tip and margin burn of Irish potatoes, 141; technique for feeding, 246–247
Intracellular bodies, occurrence of, in spike disease of sandal and Vinca, 815–817
Investigations of Aspergilli by serological methods, 148
Iris versicolor, susceptibility to Bacillus carotovorus, B. phytophthorus, and B. aroideae, 833–835
Irrigation as a cause of white spot of alfalfa, 136–137
Isolation of single spores, apparatus for, 1017–1021; method of, 17–19

JAGGER, IVAN C., The brown-blight disease of lettuce, 949–950
Jasminum primulinum, 952
JEHLE, R. A., and H. A. HUNTER, Observations on the discharge of ascospores of Venturia inaequalis in Maryland, 943–945
JOHANN, HELEN, Grated carrot agar favorable for studies of Pythium, 710; Penicillium injury to corn seedlings, 239–242
JOHNSON, E. M., see VALLEAU, W. D.
JOHNSON, H. W., see PETERSON, P. D.
JOHNSON, JAMES, Constant-temperature and humidity chambers, 227–238; Further studies on the attenuation of plant viruses, 156; The properties and behavior of potato rugose mosaic, 141
JOHNSON, T., The occurrence of yellow stripe rust in western Canada, 481; see NEWTON, M.
JONES, J. P., The influence of cropping systems and fertilizers on black and brown root rot of tobacco, 131
JONES, L. R., and REGINA S. RIKER, Studies upon the Fusarium wilt of China aster, 150
Juglans, californica, 948–949; cinerea, lesions produced by Gnomonia leptostyla, 746; hindsii, 948–949

KEITT, G. W., and E. E. WILSON, Fall applications of fungicides in relation to apple-scab control, 146; see BENSAUDE, M.; see HAMILTON, J. M.; see RIKER, A. J.
KLOTZ, L. J., Inhibition of enzymatic action as a possible factor in the resistance of plants to disease, 953
Knots, callus, on apple, grafting as a further means of preventing, 127–128
Knotweed, Uromyces polygoni on, experimental control of teliospore and urediniospore formation, 178–179, 206
KOHL, EDWIN J., The cycle of infection in apple blotch, 145
Krankheiten, Die, der Obstbäume und Obststräucher, by E. EWERT, book review, 715
KUNKEL, L. O., Further studies on the host range of aster yellows, 156

LACKEY, C. F., see CARSNER, EUBANKS
LEE, H. ATHERTON, and W. DWIGHT PIERCE, Bacterial red-stripe disease of sugar cane in countries of the Pacific, 945
LAMBERT, E. B., see E. C. STAKMAN
Lathyrus odoratus, susceptibility to Erostrotheca multiformis, 839–844
LEACH, L. D., see S. M. DIETZ
Leaf-hopper injury of legumes, 137–138
Leaf hoppers, transmission of sugar-beet curly top by, 675–679, 681–690
Leaf spot, angular, and fruit rot of cucumber, 133; ascomycetous, of cowpea, 701–704; new, of pecan, preliminary report on, 133–134
Leaf-spotting fungi, histologic changes induced in leaves, 717–751
Leaves, histologic changes induced in, by leaf-spotting fungi, 717–751
LEE, H. ATHERTON, and J. P. MARTIN, Oxidizing agents in sulphur to increase fungicidal activity, 1026–1027; and J. S. MCHARGUE, The effect of a manganese deficiency on the sugar-cane plant and its relationship to Pahala blight of sugar cane, 775–786
Legumes, leaf-hopper injury to, 137–138
LEPIK, E., Differential staining of Peronosporaceae, 869–872
Leptosphaerulina vignae, n. sp., cause of leaf spot of cowpea, 701–704
Lesions, cicatrice definitely formed about margin of, 723–731; cicatrice not definitely formed about edge of, 731–747
Leskaceae, 952
Lettuce, brown blight of, 949–950
Leucolepis acanthoneura, 952
LEVINE, MICHAEL, Studies on crown-gall transplants, 130
LEVINE, MOSES N., Biometrical studies on the variation of physiologic forms of Puccinia graminis tritici and the effects of ecological factors on the susceptibility of wheat varieties, 7–123
Light, monochromatic, photography in the study of mosaic diseases, 154; relation to experimental control of teliospore and urediniospore formation, 163–180, 206–207
Lilium auratum, mechanical transmission of mosaic of, 1025–1026
Lime, effects of, on cigar tobacco, 130–131; sulphur, fungicidal action, 355–360
Linum, reaction to rust and powdery mildew, 478
Locustidae, transmission of potato spindle tuber by, 445–448; transmitting potato spindle tuber, 140
Longevity of Pseudomonas tumefaciens Sm. & Town. in various soils, 129
Losses, determination of, in selfed lines of smutted corn, 599–602
Lycopersicum esculentum, lesions produced by Alternaria solani, 740

MACKIE, J. R., Localization of resistance to powdery mildew in the barley plant, 901–910
MACKIE, W. W., A field method of insuring positive attack with some cereal diseases, 617–621; Inheritance of resistance to blast in oats, 948
Macrosporium cucumerinum, lesions produced on Cucumis melo, 740–741
Maggot, seed-corn, transmission of potato black leg by Hylemyia cilicrura in Maine, 459
MAINS, E. B., Inheritance of resistance to Puccinia sorghi in maize, 138; and DOROTHY THOMPSON, Studies on snapdragon rust, Puccinia antirrhini, 150
Maize, inheritance of resistance in, to Puccinia sorghi, 138; root rot caused Pythium arrehomanes, 873–875; see corn
MAJOR, T. G., see SLAGG, C. M.
Manganese deficiency, relationship to Pahala blight of sugar cane, 775–786
Manihot esculenta, mosaic of, in West Africa, 155
MARCHAL, E., First report of the occurrence of Black scab or warty disease of potato in Belgium, 126
Margin, burn, of Irish potatoes, relationship of insect carriers to development of, 141
MARLOTH, R. H., see DAVIS, A. R.
MARTIN, G. HAMILTON, and VERA K. CHARLES, Preliminary studies of the life

history of Erostrotheca multiformis, the perfect stage of Cladosporium album Dowson, 839–846

MARTIN, J. P., see LEE, H. ATHERTON

MASSEY, L. M., Dry rot of Gladiolus corms, 519–529

MATSUMOTO, TAKASHI, Preliminary note on some serological studies of Aspergilli, 691–696; Some investigations of Aspergilli by serological methods, 148

McCUBBIN, W. A., Permanent mounts of white-pine-blister-rust aecia, 1029

McCULLOCH, LUCIA, and CHARLES THOM, A corm rot of Gladiolus caused by a Penicillium, 151; **Bacterium maculicola** (McC.), nom. emend. syn. Bact. maculicolum, 460; **Bact. stizolobii** (Wolf), comb. nov. syn. Aplanobacter stizolobii, 460

McHARGUE, J. S., and LEE, H. ATHERTON

McKINNEY, H. H., Virus diseases observed by the Allison V. Armour Expedition, 155 Measurements of total sunlight intensity with reference to the ecology of plant diseases, 951

Mechanical transmission of mosaic of Lilium auratum, 1025–1026

Medicago lupulina, lesions produced by Pseudopeziza medicaginis, 752; sativa, lesions produced by Pseudopeziza medicaginis, 741–742

Melampsora humboldtiana, on Salix species, 705–708; lini, reaction of Linum species to, 478

Melampsorella elatina, on Cerastium and Stellaria species, 705–708

MELHUS, I. E., and D. R. PORTER, The use of selection and selfing in improving Iacope cabbage, 142; J. H. MUNCIE, and VERNON C. FISK, Grafting as a further means of preventing callus knots on apple, 127–128

Menispermum canadense, lesions produced by Cercospora menispermi, 736–737

Mercuric, chloride and cyanide, in fire-blight-eradication work, 710–711; cyanide and chloride, in fire-blight-eradication work, 710–711

Mercury as a control for turf diseases, 137

Method, field, of epidemic induction of cereal disease, 617–621; of measuring the relative efficiency of seed-corn disinfectants, 140

Methods, for eradicating species of Rhamnus, 138; isolation of single spores, 17–19; permanent mounts of white-pine-blister-rust aecia, 1029; serological, investigations of Aspergilli by, 148; spore measurements, 19–20; technique for feeding homopterous insects, 246–247; uniform wheat rust nurseries, 13–17

Methyl violet for staining Bacillus tracheiphilus, 803–807

Microsphaera alni, 904

Migration of Bacterium tumefaciens, in tomato, 129; in tomato tissue, 553–564

Mildew, powdery, control of, 847–860; powdery, in barley plant, 901–910; powdery, of raspberry, 787–796; powdery, parasitizing Chinese cabbage, 611–615

Milesia, kriegerina, on Dryopteris spinulosa, 705–708; marginalis, on D. marginalis, 705–708; polystichi, on polystichum munitum, 705–708; pyenograndis, on Polypodium vulgare, 705–708

Millet, control of smut, 479

Mniaceae, 952

Moisture, relation to development of aseigerous stage of Venturia inaequalis, 409–414; relation to experimental control of teliospore and urediniospore formation, 181–184, 207–208; soil, effect of, on mycorrhizae, 257; soil, effect of, on sugar-beet root rot, 273–275

Monochromatic-light photography in the study of mosaic diseases, 154

MONTEITH, JOHN, JR., Leaf-hopper injury of legumes, 137–138; and A. S. DAHL, Mercury as a control for turf diseases, 137

MONTREUIL, J. E., see SLAGG, C. M.

Morphological and other characters in relation to stem-rust resistance in mature wheat plants, 650–656

Morphology (morphometry) of some important physiologic forms of Puccinia graminis tritici, 40–81; of Cladosporium strains, 324–325

Mosaic, and streak, of tomato, soil transmission of, 155; cucumber, resistance of Chinese cucumber to, 143; cytological

studies of plant tissues affected with, 154; disease of Gladiolus, 215–217; diseases, studied with monochromatic light, 154; effects upon chlorophyll content of tobacco, 697–700; of bean, transmission of, 155; of cucumber, transmitted by aphis, 143; of Lilium auratum, mechanical transmission of, 1025–1028; potato rugose, properties and behavior of, 141; tobacco, true, experiments and observations on the control of, 132; virus, a "streak" of tomatoes produced by disturbing principle from healthy potatoes in combination with, 311; virus, of tobacco, accuracy in comparing various concentrations of, 132

Moss, E. M., The haustorium of Cuscuta gronovii, 478

Mottle-leaf disease of beech, 151–152

Mounts, permanent, of white-pine-blister-rust aecia, 1029

Muncie, J. H., Rumex crispus, a weed host of Pseudomonos tumefaciens, 130; see Melhus, I. E.; and W. B. Shippy, Overgrowths and hairy root on nursery-apple and quince trees, 127

Mycorrhizae, sugar-cane, and their relation to root diseases, 249–261

Mycorrhizal fungus of Vaccinium, 148

Mycosphaerella, fragariae, lesions produced on Fragaria virginiana, 774; grossulariae, lesions produced on Ribes prostratum, 744–745; sentina, lesions produced on Pyrus communis, 727

Narasimhan, M. J., Note on the occurrence of intracellular bodies in spike disease of sandal (Santalum album Linn.), 815–817

Neal, D. C., Outstanding wilt-resistant-cotton varieties, 134

Neatby, K. W., see Goulden, C. H.

Needle rusts on balsam firs in North America, conspectus of, 705–708

Nematodes, cause of root knot of abacá, 861–867

Nemopanthus mucronata, lesions produced by Rhytisma ilices-canadensis, 743–744

Newton, George A., Some fungi of the Stemphylium type and their relation to apple rots, 565–578

Newton, M., see Brown, A. M.; see Johnson, T.; T. Johnson, and A. M. Brown, Physiologic forms of wheat stem rust in Canada, 478–479

Nicotiana glauca, mosaic of, on Canary Islands, 155

Nisikado, Y., Review of Shirai and Hara's book, A list of Japanese fungi hitherto unknown, 1023

Notes, phytopathological, 125–126, 245–248, 311–312, 459–461, 1025–1030; Ericksson prizes, 1029–1030; field meeting, Southern Section, American Phytopathological Society, 460–461.

Nuckols, S. B., and Tompkins, C. M.

Nurseries, uniform wheat-rust, 13–17, 20–31

Nursery trees, apple and quince, overgrowths and hairy root on, 126

Nutrients, soil, effect on mycorrhizae, 257–258

Oat, blast, 948; dust treatment for smuts of, 139

Oats, Fusarium blight of, 480; physiologic forms of stem rust in Canada, 479–480; varietal resistance to Ustilago levis and U. avenae, 993–997

Observations on the discharge of ascospores of Venturia inaequalis in Maryland, 943–945

Obstbäume und Obststräucher, Die Krankheiten der, book review, 715

Occurrence, of intracellular bodies in spike disease of sandal (Santalum album), 815–817; of Peronospora sparsa Berk. on hothouse roses in southern California, 950; of pycnia and aecia in certain rust fungi, 1005–1015

Ocfemia, Gerardo Offimaria, and Melanio R. Calinisan, The root knot of abacá, or Manila hemp, 861–867

O'Gara, Patrick Joseph, In memoriam, 475

Ogilvie, Lawrence, "Black tip," a finger-tip disease of the Chinese banana in Bermuda, 531–538

Oidium sp., relation of Linum species to, 478

Oleander bacteriosis, 503–518

O'Leary, Keith, Eocronartium musicola

fr. in a new locality and on a new host, 952
Olives, copper sulphate as remedy for exanthema in, 449–454
Onoclea sensibilis, host of Uredinopsis mirabilis, 705–708
Oospora citri-aurantii, 949
Ophiobolus graminis Sacc., the reaction of wheat varieties to inoculations with, 477
Origin of Phytophthora infestans, 494–500
Ornamental plants, condition of bulbs and corms of, offered for importation into Canada, 150–151
ORTON, C. R., The effect of disinfectants upon the germination of seeds kept in storage for indefinite periods after treatment, 136
Osmorhiza longistylis, lesions produced by Septoria osmorhizae, 734–735
Osmunda species, hosts of Uredinopsis osmundae, 705–708
Overgrowth, wound, correlation of, and crown gall of apple, 128
Overgrowths, and hairy root on nursery-apple and quince trees, 127; wound, studies of the history of development of, and of the influence of wrappers on their suppression, 128
Oxidizing agents in sulphur to increase fungicidal activity, 1026–1027

Pahala blight of sugar cane, effect of manganese deficiency on, 775–786
Palm, Washingtonia, leaf spot due to Cylindrocladium macrosporum, 219–225
PAMMEL, L. H., Cladosporium diseases, 946
Paragynous antheridia of Phytophthora spp., 149–150
Parasitism, (host plants, host range, pathogenicity), tobacco ring spot, 133; host range of aster yellows, 156; of Aphanomyces euteiches on conifers, 366; of Bacilus spp. to potato tubers, 819–838; of Corticium koleroga, 147–148; of physiologic forms of cereal smuts, 981–997; of physiologic forms of Tilletia levis and T. tritici, 985–992; of physiologic forms of Ustilago hordei, 992–993; of Pythium spp. on conifers, 363–366; relation of variety of apple to production of ascigerous stage of Venturia inaequalis, 393–395
Paratrioza cockerelli, its relation to a new potato disease in Utah, 140–141
PATEL, M. K., Longevity of Pseudomonas tumefaciens Sm. & Town. in various soils, 129; Strains of Ps. tumefaciens Sm. & Town. and their prevalence in various soils, 129–130; A study of pathogenic and nonpathogenic strains of tumefaciens Sm. & Town., 331–343
Pathogenicity, of Fusarium vasinfectum, 419–438; of Sclerotium gladioli, 524
PATTERSON, FLORA W., 877–879
Peach scab, caused by Cladosporium, 313
Pears, copper sulphate as remedy for exanthema in, 499–454
Pecan, preliminary report of a new leaf spot of, 133–134
Pelargonium zonale, susceptibility to Bacillus carotovorus, B. aroideae, and B. phytophthorus, 834–835
PELTIER, GEORGE L., Control of bunt of wheat in Nebraska, 921–929
Penetration, mode of, by Cladosporium carpophilum, 316–318; of furfural in plant tissues, 152
Penicillium, causing a corm rot of Gladiolus, 151; corm rot of Gladioli, 151; digitatum, 949; Gladioli, corm rot of Gladioli, 151; injury to corn seedlings, 239–242; italicum, 949; oxalicum, possible cause of injury to corn seedlings, 239–242
Pentstemon, rust on, 665
Peony, the crown-elongation disease of, 243–244; Phytophthora blight of, 881–900
Peridermium, balsameum, on Abies species, 705–708; columnare, on Abies species, 705–708; elatinum, on Abies species, 705–708; kriegerina, on Abies balsamea, 705–108; marginalis, on Abies balsamea, 705–108; ornamentale, on Abies species, 705–708; pustulatum, on Abies species, 705–708, pycnoconspicuum, on Abies balsamea, 705–708; pycnogrande, on Abies balsamea, 705–708; rugosum, on Abies species, 705–708; sp., possible

aecial stage of Puccíniastrum myrtilli, 623–625
Perithecia of Venturia inaequalis, factors important in development of, 145–146
Permanent mounts of white-pine-blister-rust aecia, 1029
Peronospora sparsa, 950
Peronosporaceae, differential staining of, 869–872
PETERSON, P. D., and H. W. JOHNSON, Powdery mildew of raspberry, 787–796
Petri-dish experiments for control of teliospore and urediniospore formation, 184–199
Phaseolus vulgaris, Uromyces appendiculatus on, experimental control of teliospore and urediniospore formation, 163–169, 183, 184–187, 206
Phegopteris dryopteris, host of Hyalospora aspidiotus, 705–708; host of Uredinopsis phegopteridis, 705–708
Phoma, 917; betae, 941; sp., 952
Phomopsis, californica, 949; sp., causing cranberry storage rot, 809–814
Photographic collection of ERWIN F. SMITH, 312
Photography, monochromatic-light, in the study of mosaic diseases, 154
Phyllosticta fraxinii, lesions produced on Fraxinus americana, 741
Phymatotrichum (Ozonium) omnivorum, 952–953
Physiologic specialization, biometrical studies of physiologic forms of Puccinia graminis tritici, 40–81; cultural differences between physiologic forms of cereal smuts, 958–972; effect of temperature on growth of physiologic forms of cereal smuts, 972–981; forms of P. graminis avenae in Canada, 479–480; forms of wheat stem rust in Canada, 478–479; in P. sorghi, 349–352; in some cereal smuts, 955–1003; new forms of Tilletia levis and T. tritici, 579–588; new physiologic forms of T. tritici in wheat, 139; pathogenic differences between physiologic forms of cereal smuts, 981–997; stability of physiologic forms of P. graminis tritici under diverse cultural conditions, 108–113; strains of Cladosporium affecting stone fruits, 318–322, 327–328; strains of Pseudomonas tumefaciens, 129–130; two forms of Ustilago striaeformis (Westd.) Neissl., 149
Phytomonas malvacera, 951
Phytopathological Society, see American Phytopathological Society
Phytophthora, 881–882, 884–886, 890–895, 897; blight, 881–882, 896–897; cactorum, 881, 884, 889, 894–896, 898, 948–949; colocasiae, 894; cryptogea, 894; erythroseptica, 893–895, 898; faberi, 895; fagi, 881, 894–895; in relation to crown rot of walnut, 948–949; infestans, 894–895; infestans, origin of, 494–500; jatrophae, 895; mexicana, 894; nicotianae, 889–890, 895; omnivora, 895; paeoniae, 881, 896–899; palmivora, 894; parasitica, 894–895; phaseoli, 895; pipi, 894; richardiae, 895; rot of apple, 885; rot of ginseng, 885; rot of strawberry, 885; species as stock cultures, 247–248; spp., paragynous antheridia of, 149–150; syringae, 895; terrestris, 894–895
PIERCE, W. DWIGHT, see LEE, H. ATHERTON
Pima cotton, 951
Pine, white, permanent mounts of blister-rust aecia of, 1029
Pink rot, 931
PITTMAN, D. W., see STEWART, GEORGE
PLAKIDAS, A. G., Strawberry dwarf, 439–444
Plants, disease-resistant, cytological phenomena in, 144
Plasmodiophora, brassicae, in relation to soil alkalinity, 301–306; reaction and treatment of soils infested with, 141–142
Plasmopara viticola, differential staining of, 869–872
Pleospora mali, n. sp., cause of apple rot, 576
Plum, scab caused by Cladosporium, 313–314
PLUNKETT, O. A., The occurrence of Peronospora sparsa Berk. on hothouse roses in southern California, 950
Podophyllum peltatum, lesions produced by Septoria podophyllina, 731–732
Polygala senega, rust on, 665

Polygonum aviculare, Uromyces polygoni on, experimental control of teliospore and urediniospore formation, 178–179, 206
Polypodium vulgare, host of Milesia pycnograndis, 705–708
Polystichum munitum, host of Milesia polystichi, 705–708
POOLE, R. F., Sweet-potato stem rot prevented by treating stems and roots with Bordeaux mixture, 152–153
Populus sp., 952
PORTER, C. L., see COOPER, D. C.
PORTER, D. R., see MELHUS, I. E.; Studies with the watermelon wilt, caused by Fusarium niveum E. F. Sm., 143–144; Varietal resistance of watermelon to wilt (Fusarium niveum E. F. Sm.), 144
PORTER, R. H., Further evidence of resistance to cucumber mosaic in the Chinese cucumber, 143; Seed disinfectants for the control of covered smut and stripe of hulless barley, 139; T. F. YU and H. K. CHEN, The effect of seed disinfectants on smut and yield of millet, 911–919
Potato, a new and destructive disease of, in Utah, 140–141; black scab or warty disease of, in Belgium, 126; Irish, relationship of insect carriers to development of tip and margin burn of, 141; pathogenicity of Bacillus mesentericus, B. aroideae, B. carotovorus, and B. phytophthorus to, 819–832; psylla of, its relation to a new potato disease in Utah, 140–141; rugose mosaic of, properties and behavior of, 141; transmission of black leg by seed-corn maggot, 459; transmission of spindle tuber by grasshoppers, 445–448; transmission of spindle tuber of, 140
Potatoes, blight-resistant, 483–502; varietal susceptibility of, to Fusarium wilt and stem-end rot, 307–309
Powdery mildew, and rust, reaction of Linum species to, 478; control of, 847–860; of raspberry, 787–796
Pox, or soil rot, in sweet potato, caused by an Actinomycete, 152
Predisposition, condition of host plants to severity of stem-rust epidemics, 83–90; of sugar beets to late root rot, 263–276
Preliminary note on some serological studies of Aspergilli, 691–696
Prenanthes alba, Puccinia orbicula on, experimental control of teliospore and urediniospore formation, 197–199
Primula polyantha, lesions produced by Ramularia primulae, 738
Prize offered by the State of Rio de Janeiro, Brazil, 713
Prizes, Eriksson, announcement of, 1029–1030
Production of agglutinins by phytopathogenic bacteria, 277–288
Progress of rust studies, 659–674
Properties and behavior of potato rugose mosaic, 141
Prunes, copper sulphate as remedy for exanthema in, 449–454
Prunus, armeniaca, 952; avium, lesions produced by Coccomyces hiemalis, 727; cerasus, lesions produced by C. hiemalis, 726–727; domestica, lesions produced by C. prunophorae, 723–725; serotina, lesions produced by Cercospora circumcissa, 737–738; virginiana, lesions produced by Coccomyces lutescens, 725–726
Pseudomonas, cerasus, production of antisera against, 278–288; savastanoi var. nerii, cultural characteristics, 513–515; savastanoi var. nerii, host relationship, 512–513; tumefaciens, a study of pathogenic and nonpathogenic strains, 331–343; tumefaciens, on Rumex crispus, a weed host of, 130; tumefaciens, see Bacterium tumefaciens; tumefaciens Sm. & Town, longevity of, in various soils, 129; tumefaciens Sm. & Town., strains of, and their prevalence in various soils, 129–130
Pseudopeziza, medicaginis, lesions produced on Medicago lupulina, 472; medicaginis, lesions produced on M. sativa, 741–742; ribis, lesions produced on Ribes grossularia, 742–743; trifolii, lesions produced on Trifolium pratense, 742
Psoralea bituminosa, mosaic of, on Canary Islands, 155
Psylla, potato, its relation to a new potato disease in Utah, 140
Pteridium aquilinum and P. pubescens,

hosts of Uredinopsis macrosperma, 705–708

Puccinia, andropogonis, 665; anomala, in western Canada, 481; anomala, method of epidemic induction, 617–621; antirrhini on Antirrhinum majus, experimental control of teliospore and urediniospore formation, 176–178, 183–184, 192–194; antirrhini, rust of snapdragon, 150; asparagi on Asparagus officinale, experimental control of teliospore and urediniospore formation, 180, 181–183; comandrae, 665; coronata, heterothallism in, 1005–1015; coronata, method of epidemic induction, 617–621; coronata, on species of Rhamnus, 138; glumarum, in western Canada, 481; graminis avenae, physiologic forms in Canada, 479–480; graminis, heterothallism in, 147, 1005–1015; graminis, method of epidemic induction, 617–621; graminis, rust epidemic induction, 617–621; graminis, rust epidemics and temperature, 372–374; graminis, sexual behavior of, 479; graminis tritici, biometrical studies on the variation of physiologic forms of, and the effects of ecological factors on the susceptibility of wheat varieties to, 7–123; graminis tritici, consistency in behavior of different strains of identical forms of, 112–113; graminis tritici, field observations on severity of epidemics, 20–31; graminis tritici, greenhouse and laboratory studies of physiologic forms of, 31–81; graminis tritici, identity and nature of physiologic forms isolated from uniform rust nurseries in 1919–1923, 33–35; graminis tritici, inheritance of resistance to, in a cross between two varieties of Triticum vulgare, 631–658; graminis tritici, inoculation technique and cultural methods in greenhouse, 17–20; graminis tritici, morphology of some important physiologic forms of, 40–81; graminis tritici, physiologic forms in Canada, 478–479; graminis tritici, prevalence and distribution of physiologic forms isolated from uniform rust nurseries in 1919–1923, 35–40; graminis tritici, single-spore isolations and spore measurements of, 17–20; graminis tritici, stability of physiologic forms under diverse cultural conditions, 108–113; graminis tritici, uniform rust nurseries, 13–17, 20–31; graminis tritici, uniformity of behavior over long periods of time of different physiologic forms of, 108–112; helianthi, heterothallism in, 147, 1005–1015; orbicula on Prenanthes alba, experimental control of teliospore and urediniospore formation, 197–199; pentstemonis, 665; podophylli, 671; pringsheimiana, heterothallism in, 1005–1015; rubigo-vera, 666–667; seymeriae, 665; sorghi, inheritance of resistance to, in maize, 138; sorghi, on Zea mays, experimental control of teliospore and urediniospore formation, 179–180, 194–195, 206; sorghi, physiologic specialization, 345–354; suaveolens on Cirsium arvense, experimental control of teliospore and urediniospore formation, 169–172, 191–192, 206; taraxaci on Taraxacum officinale, experimental control of teliospore and urediniospore formation, 175–176, 196–197; triticina, method of epidemic induction, 617–621; triticina on Triticum vulgare, experimental control of teliospore and urediniospore formation, 179, 195–196

Pucciniastrum, abietichamaenerii, on Epilobium species, 705–708; epilobii, on Epilobium species, 705–708; myrtilli, on Vaccinium species, 705–708; myrtilli, possible alternate stage of, 623–625

Pycnia, occurrence of, in certain rust fungi, 1005–1015; occurrence under natural conditions, unaccompanied by aecia, 1010–1014

Pyrus communis, lesions produced by Mycosphaerella sentina, 727

Pythiacystis, 953; citrophthora, 949; species as stock cultures, 247–248

Pythium, arrhenomanes, cause of maize root rot, 873–875; arrhenomanes, comparison with P. aphanidermatum, 874; grated-carrot agar for, 710; species as stock cultures, 247–248; spp., pathogenicity for conifers, 363–366

QUANJER, H. M., book review, Die Krankheiten der Obstbäume und Obststräucher,

by E. EWERT, 715; Eriksson prizes, 1029–1030
Quercus pedunculata, 904
Quince, and apple, nursery trees, overgrowths and hairy root on, 127

RALEIGH, W. P., A preliminary method of measuring the relative efficiency of seed-corn dinifectants, 140
Ramularia, **carniformis** (E. & T.), n. com. on Tripsacum dactyloides, 149; decipiens, lesions produced on Rumex crispus, 739; primulae, lesions produced on Primula polyantha, 738; ranunculi, lesions produced on Ranunculus recurvatus, 738–739
RANKIN, W. HOWARD, Mottle-leaf disease of beech, 151–152
Ranunculus recurvatus, lesions produced by Ramularia ranunculi, 738–739
Raspberry, powdery mildew, 787–796
Reaction and treatment of soils infested with Plasmodiophora brassicae Wor., 141–142
Red stripe, bacterial, disease of sugar cane, 945
REDDICK, DONALD, Blight-resistant potatoes, 483–502
Relationship of insect carriers to the development of tip and margin burn of Irish potatoes, 141
Report, of nineteenth annual meeting of American Phytopathological Society, 463–474; of ninth annual meeting of Canadian Division of American Phytopathological Society, 477–481; of progress, on condition of bulbs and corms of ornamental plants offered for importation into Canada, 150–151; on a new leaf spot of pecan, 133–134; on insect carriers of tip and margin burn of Irish potatoes, 141; on mechanical transmission of mosaic of Lilium auratum, 1025–1026
Resistance, cytological phenomena in disease-resistant plants, 144; Ekishirazu, blight-resistant variety of potato, 484; inheritance of, to Puccinia sorghi in maize, 138; of host, relation to teliospore formation, 208–209; of varieties of watermelons to wilt (Fusarium niveum), 144; relative, of selfed lines of corn to rust, 345–349; see susceptibility; to cucumber mosaic in Chinese cucumber, 143; to Puccinia graminis tritici, inheritance of, in a cross between two varieties of Triticum vulgare, 631–658; to stem rust in mature plants in relation to morphological and other characters, 650–656; varietal, of cereal crops to physiologic forms of smut, 981–997
Review, of A list of Japanese fungi hitherto unknown, 1023; of Plantsiektes hul oorsaak destryding, 1023
Rhabdospora oxydendri, (E. & E.), n. com. on Oxydendrum arboreum, 149
Rhamnus, effective methods for eradicating species of, susceptible to Puccinia coronata, 138
Rhynchosporium secalis, method of epidemic induction, 617–621
Rhytisma, acerinum, lesions produced on Acer rubrum, 744; ilices-canadensis, lesions produced on Ilex verticillata, 743; ilices-canadensis, lesions produced on Nemopanthus mucronata, 743–744
Ribes, grossularia, lesions produced by Pseudopeziza ribis, 742–743; prostratum, lesions produced by Mycosphaerella grossulariae, 744–745
RICHARDS, B. L., A new and destructive disease of the potato in Utah and its relation to the potato psylla, 140–141; Irrigation as a cause of white spot of alfalfa, 136–137
RIKER, A. J., A bacterial disease of broad beans, 136; Correlation of the wound overgrowth and crown gall of apple in parts of Europe and of the United States, 128; Notes on crown gall situa- in England, France and Holland, 289–294; W. M. BANFIELD and G. W. KEITT, Studies of the history of development of wound overgrowths on apple grafts and of the influence of wrappers on their suppression, 128
RIKER, REGINA S., see JONES, L. R.
Ring spot of tobacco, a virus disease with a wide host range, 133
RODENHISER, H. A., Experiments on the control of barley stripe, 295–300; Physio-

logic specialization in some cereal smuts, 955–1003

Root, and crown injury, apple trees, 547–551; club, in relation to soil alkalinity, 301–306; diseases, observations on sugar-cane mycorrhizae and their relation to, 249–261; diseases, prevention of sugar cane, 259; knot, of abacá, 861–867; knot, symptoms on abacá, 861–863; rot, black and brown, of tobacco, influence of cropping systems and fertilizers on, 131; rot, black, of apple, 145; rot, black, of tobacco and soil reaction, 131; rot, brown, of tobacco, effect of timothy infusions on, 131–132; rot, late, predisposition of sugar beets to, 263–276; rot, Texas, 952–953

Rosen, H. R., A consideration of the pathogenicity of the cotton-wilt fungus, Fusarium vasinfectum, 419–438; A preliminary report on the relationship of insect carriers to the development of tip and margin burn of Irish potatoes, 141

Roses, occurrence of Peronospora sparsa Berk. on, 950

Rot, black, of sweet potatoes, treatment for control of, 153–154; corm, of Gladiolus, caused by a Penicillium, 151; dry, of Gladiolus corms, 519–529; in sweet potato, caused by an Actinomycete, 152; Penicillium, corm, of Gladioli, 151; root, caused by Pythium arrhenomanes, 873–875; stem-end, varietal susceptibility of potatoes to Fusarium wilt and, 307–309

Rots, apple, relation of Stemphylium type of fungi to, 565–578; storage, of cranberries, in 1927, 809–814

Rumex crispus, lesions produced by Ramularia decipiens, 739; a weed host of Pseudomonas tumefaciens, 130

Russell, R. C., The reaction of wheat varieties to inoculations with Ophiobolus graminis Sacc., 477

Rust, and powdery mildew, reaction of Linum species to, 478; blister, of white pine, permanent mounts of aecia of, 1029; epidemics, wheat stem, 369–374; fungi, heterothallism in, 147; fungi, occurrence of pycnia and aecia in, 1005–1015; fungi, teliospore and urediniospore formation controlled experimentally, 157–213; inheritance of resistance in mature plants in relation to morphological and other characters, 650–656; leaf and stem, sulphur dusting for control of, 480; leaf, dwarf, on barley in western Canada, 481; of corn, Puccinia sorghi, 345–354; of snapdragon, P. antirrhini, 150; of wheat, effect of ecological factors on susceptibility of varieties to, 81–108; stem, inheritance of resistance to, in wheat, 631–658; stem, of wheat, see P. graminis tritici; stem, physiologic forms of, on oats, in Canada, 479–480; stem, physiologic forms of, on wheat, in Canada, 478–479; stripe, occurrence in western Canada, 481; studies, progress of, 659–674

Rusts, needle, on balsam firs in North America, conspectus of, 705–708; of the Pacific Northwest, 948

Salix species, alternate hosts of fir-needle rust, 705–708

Sandal, occurrence of intracellular bodies in spike diesease of, 815–817

Sanford, G. B., Cereal diseases in Alberta in 1927, 481

Santalum album, occurrence of intracellular bodies in spike disease of, 815–817

Savastano, G., and H. S. Fawcett, The effect of mixed inoculations of certain citrus fruit-rotting organisms, 949

Sayre, J. D., and R. C. Thomas, New dust treatments for oat smuts, 139

Scab, apple, see Venturia inaequalis; black, or warty disease of potato, in Belgium, 126; of apple, certain sulphur fungicides in control of, 146–147; of apple, fall applications of fungicides in relation to control of, 146

Scandiffio, Mario, see Church, Margaret B.

Scheffer, T. C., Steam sterilization of coniferous seed beds, 952

Sclerospora graminicola, 917

Sclerotinia libertiana, 949

Sclerotium gladioli, n. sp., etiology of dry rot of Gladiolus, 521–524, pathogenicity.

524, cultural characters, 524–526, life history, 526–527, control, 527–528
Seed, cotton, treatment by the dusting method, 134; disinfectants, effect on smut and yield of millet, 911–919; disinfectants for control of covered smut and stripe of hulless barley, 139; effect of washing on infection of wheat by stinking smut, 711–713
Seedling blight, on oats, caused by Fusarium culmorum, 480
Seeds, effect of disinfectants upon germination of, in storage, 136
Selection and selfing for improving Iacope cabbage, 142
Selfing for improving Iacope cabbage, 142
Septoria, acerina, lesions produced on Acer pennsylvanicum, 734; cirsii, lesions produced on Cirsium arvense, 733; conspicua, lesions produced on Steironema ciliatum, 732–733; of tomato, control by means of forced ventilation in hotbeds, 1027–1029; osmorhizae, lesions produced on Osmorhiza longistylis, 734–735; petroselini, lesions produced on Apium graveolens, 731; podophyllina, lesions produced on Podophyllum peltatum, 731–732; tritici, method of epidemic induction, 617–621; verbasicola, lesions produced on Verbascum blattaria, 733–734
Serological, investigations of Aspergilli, 148; studies on Aspergilli, 691–696
Setaria italica, 911
SEVERIN, HENRY H. P., Tomato yellows or tomato curly top, 709–710; and OLIVE SWEZY, Filtration experiments on curly top of sugar beets, 681–690
Sex in Puccinia graminis, 479
SHAPOVALOV, MICHAEL, and F. S. BEECHER, The development of tomato yellows under light conditions, 950
SHERBAKOFF, C. D., An examination of Fusaria in the herbarium of the Pathological Collections, Bureau of Plant Industry, U. S. Department of Agriculture, 148–149; Washingtonia palm-leaf spot due to **Cylindrocladium macrosporum,** n. sp., 219–225
SHIPPY, W. B., see MUNCIE, J. H.
SHIRAI, M., and K. HARA, A list of Japanese fungi hitherto unknown, review of, 1023
SIMMONDS, P. M., A seedling-blight disease of oats caused by Fusarium culmorum, 480
Single spores, see spores, single
SKORIC, VLADIMIR, see WRIGHT, WILLIAM H.
SLAGG, C. M., J. E. MONTREUIL, and T. G. MAJOR, The effects of lime on cigar tobacco, 130–131
SMITH, CLAYTON O., Oleander bacteriosis in California, 503–518; A study of citrus blast and some allied organisms, 952
SMITH, ERWIN FRINK, In memoriam, 475; Photographic collection of, 312; Testimonial to, 1–5
SMITH, RALPH E., and HAROLD E. THOMAS, Copper sulphate as a remedy for exanthema in prunes, apples, pears, and olives, 449–454
Smut, covered, of hulless barley, control of, 139; influence of environmental factors on prevalence in corn, 589–598; losses in selfed lines of corn, 599–602; oat, dust treatments for, 139; of millet, 911–919; stinking, of wheat, effect of washing seed on infection, 711–713; treatment of millet seed for prevention of, 479
Smuts, cereal, physiologic specialization in, 955–1003
Snapdragon, Puccinia antirrhini on, experimental control of teliospore and urediniospore formation, 176–178, 183–184, 192–194; rust of, antirrhini, 150
Snowberry, anthracnose of, 797–801
Soft rot, 931
Soil, alkalinity in relation to clubroot, 301–306; in relation to severity of stem-rust epidemics, 90–91; productivity, effect of, on sugar-beet root rot, 268–272; reaction, and black root rot of tobacco, 131; transmission of tomato mosaic and streak in the greenhouse, 155
Soils, longevity of Pseudomonas tumefaciens Sm. & Town. in, 129; prevalence of strains of Ps. tumefaciens in, 129–130

Solanum tuberosum, lesions produced by Alternaria solani, 739–740
Solutions, sugar, for experimental control of teliospore and urediniospore formation, 184–199; transmission of virus of curly top of sugar beet through, 675–679
Sonchus oleraceus, 904
SØRENSEN, S. P. L., Emil Chr. Hansen Medal, 1928, 125
SPALTEHOLZ, R., Review of van der Bijl's book, Plantsiektes hul oorsaak en bestryding, 1023
Sphaceloma, fawcettii Jenkins, cause of lesions on Citrus leaves, 539–545; **symphoricarpi,** n. sp., causing anthracnose of Symphoricarpos albus var. laevigatus, 799–800
Sphaerella tussilaginis, lesions produced on Tussilago farfara, 745–746
Sphaerotheca humuli, causing powdery mildew of raspberry, 787–788
Spike disease of sandal (Santalum album Linn.), occurrence of intracellular bodies in, 815–817
Spinach, host of beet curly top, 709–710
Spindle tuber, of potato, transmission of, 140; transmission by grasshoppers, 445–448
Spores, measurements, method of, 19–20; single, apparatus for isolating, 1017–1021; single, method for isolating, 17–19
Sporotrichum fruit spot of apple, 145
Stability of physiologic forms of Puccinia graminis tritici under diverse cultural conditions, 108–113
Stain, Giemsa, demonstration of bacteria in plant tissues by means of, 803–807
Staining, differential, of Peronosporaceae, 869–872
STAIR, E. C., H. D. BROWN, and T. E. HIENTON, Forced ventilation as a means of controlling tomato Cladosporium and Septoria in hotbeds, 1027–1029
STAKMAN, E. C., and E. B. LAMBERT, The relation of temperature during the growing season in the spring-wheat area of the United States to the occurrence of stem-rust epidnmics, 369–374; J. J. CHRISTENSEN, and H. E. BREWBAKER, Physiologic specialization in Puccinia sorghi, 345–354
Steam sterilization of coniferous seed beds, 952
Steironema ciliatum, lesions produced by Septoria conspicua, 732–733
Stellaria species, alternate hosts of a fir-needle rust, 705–708
Stem-end rot, varietal susceptibility of potatoes to Fusarium wilt and, 307–309
Stemphylium, congestum, n. sp., cause of apple rot, 576; type of fungi, relation to apple rots, 565–578
Stem rot, of sweet potato, prevented by treating stems and roots with Bordeaux mixture, 152–153
Stem rust, see Puccinia graminis
STEVENS, NEIL E., and HENRY F. BAIN, Storage rots of cranberries in 1927 crop, 809–814
STEWART, GEORGE, and D. W. PITTMAN, Predisposition of sugar beets to late root rot, 263–276
Stinking smut, of wheat, effect of washing seed on infection, 711–713
STONE, R E., see HOWITT, J. E.; Treatment of millet seed to prevent smut, 479
Storage, effect of disinfectants upon the germination of seeds kept in, for indefinite periods after treatment, 136; rots, of cranberries, in 1927, 809–814
STOUT, G. L., see TEHON, L. R.
STOVER, W. G., Experiments with tomato streak, 154
Strains, of Pseudomonas tumefaciens Sm. & Town. and their prevalence in various soils, 129–130; see physiologic specialization
Strawberry, dwarf, 439–444; killing of roots, 245–246
Streak, of tomato, experiments with, 154; of tomato, soil transmission of, 155; of tomatoes produced by disturbing principle from apparently healthy potatoes in combination with tomato-mosaic virus, 311
STREETS, R. B., Studies of Texas root rot in Arizona, 952–953
Stripe, barley, experiments on control of, 295–300; brown, of sugar cane, caused by **Helminthosporium stenospilum,** n.

sp., 135–136; brown, of sugar cane, in Cuba, 135; disease of barley, control of, 477–478; of hulless barley, control of, 139

Struthiopteris germanica, host of Uredinopsis struthiopteridis, 705–708

Studies, of black root rot of apple, 145; of the history of development of wound overgrowths on apple grafts and of the influence of wrappers on their suppression, 128; on crown-gall transplants, 130; on snapdragon rust, Puccinia antirrhini, 150; on the attenuation of plant viruses, 156; on the host range of aster yellows, 156; on the life history of the crown-gall organism; 128–129; upon the Fusarium wilt of China aster, 150; with watermelon wilt, caused by Fusarium niveum E. F. Sm., 143–144

Study, of citrus-blast and some allied organisms, 952; of pathogenic and non-pathogenic strains of Pseudomonas tumefaciens Sm. & Town., 331–343

Sugar beet, curly top, 951; occurrence of **Aphanomyces cochlioides**, n. sp., on, 149; transmission of curly-top virus through different solutions, 675–679

Sugar beets, diseases in storage resulting from hook injury, 939–941; filtration experiments on curly top of, 681–690; predisposition of, to late root rot, 263–276

Sugar cane, brown stripe of, 761–772; brown stripe of, caused by **Helminthosporium stenospilum**, n. sp., 135–136; brown stripe of, in Cuba, 135; effect of manganese deficiency on Pahala blight, 775–786; eye spot, ocellum as cause, 757–761, 763–770; eye spot, in Cuba, 135; gummosis of, 135; helminthosporiose of, 754–758; observations on mycorrhizae and their relation to root diseases of, 249–261; three Helminthosporium diseases of, 753–774

Sugar solutions, effect on beet curly-top virus, 675–679; for experimental control of teliospore and urediniospore formation, 184–199

Sulphur, dust, for control of leaf and stem rust, field trials, 480; dust, for control of leaf and stem rust, small-plot, 480; factors affecting the fungicidal property of, 147; fungicides, in the control of apple scab, 146–147; oxidizing agents in, to increase fungicidal activity, 1026–1027

Susceptibility, of potato varieties to Bacillus mesentericus, B. carotovorus, B. phytophthorus, and B. aroideae, 819–838; of wheat varieties, effect of ecological factors on, 81–108; see resistance; varietal, of potatoes to Fusarium wilt and stem-end rot, 307–309

Sweet potato, soil rot or pox in, caused by an Actinomycete, 152; stem rot, prevented by treating stems and roots with Bordeaux mixture, 152–153

Sweet potatoes, treatment of, with various disinfectants for black-rot control, 153–154

SWEZY, OLIVE, see SEVERIN, HENRY H. P.

Symphoricarpos albus var. laevigatus, anthracnose of, 797–801

Symplocarpus foetidus, lesions produced by Cercospora symplocarpi, 736

Take-all, see Ophiobolus graminis

Taraxacum officinale, Puccinia taraxaci on, experimental control of teliospore and urediniospore formation, 175–176, 196–197

TEHON, L. R., and G. L. STOUT, An ascomycetous leaf spot of cowpea, 701–704

Teliospore formation, experimental control of, 157–213

Temperature, chambers, constant-control, 227–238; effect of, on development of potato-tuber rots, 823–827; effect on growth of physiologic forms of cereal smuts, 972–981; optimum, for germination of Puccinia glumarum, 481; relation of, to occurrence of stem-rust epidemics, 369–374; relation to development of ascigerous stage of Venturia inaequalis, 396–409; relation to experimental control of teliospore and urediniospore formation, 163–180; relations of Cladosporium, 322–324; relations, on culture media, of Bacillus aroideae, carotovorus, mesentericus, phytophthorus, 822–823

Terminology, a question of correct usage, 460
Testimonial to ERWIN FRINK SMITH, 1–5
Thistle, Puccinia suaveolens on, experimental control of teliospore and urediniospore formation, 169–172, 191–192, 206
THOM, CHARLES, see MCCULLOCH, LUCIA
THOMAS, H. E., Killing of strawberry roots, 245–246; Root and crown injury to apple trees, 547–551
THOMAS, HALOLD E., see SMITH, RALPH E.
THOMAS, R. C., see SAYRE, J. D.
THOMPSON, DOROTHY, see MAINS, E. B.
Thread blight, caused by Corticium koleroga, 147–148
Tilletia, levis, effect of washing seed on infection, 711–713; levis, new physiologic forms of, 579–588; levis, pathogenicity of physiologic forms of, 985–992; levis, varietal resistance of spring wheat to, 981–992; tritici, new physiologic forms of, 579–588; tritici, new physiologic forms of, in wheat, 139; tritici, pathogenicity of physiologic forms of, 985–992
Timothy infusion, effect of different ages of, on growth of tobacco and on brown root rot of tobacco, 131–132
Tip burn of Irish potatoes, relationship of insect carriers to development of, 141
Tissues, plant, cytological studies of, affected with mosaic diseases, 154; plant, demonstration of bacteria in, by means of Giemsa stain, 803–807; plant, differential staining of Peronosporaceae in, 869–872; plant, migration of Bacterium tumefaciens in, of tomato, 553–564; plant, penetration of furfural in, 152; plant, relation of leaf infection to production and of time of leaf fall to development of ascigerous stage of Venturia inaequalis, 375–393
Tobacco, black root rot of, and soil reaction, 131; cigar effects of lime on, 130–131; effect of timothy infusion of different ages on the growth of, 131–132; effects of mosaic upon chlorophyll content of, 697–700; experiments and observations on control of true mosaic of, 132; influence of cropping systems and fertilizers on black and brown root rot of, 131; mosaic virus, accuracy in comparing various concentrations of, 132; ring spot, 133; virus diseases of, in Kentucky, 132–133
Tomato, control of Cladosporium and Septoria of, by means of forced ventilation in hotbeds, 1027–1029; curly top, or yellows, 709–710; experiments with streak of, 154; method and rate of migration of Bacterium tumefaciens in, 129; migration of Bact. tumefaciens in tissue of, 553–564; mosaic and streak of, soil transmission, 155; susceptibility to Bacillus aroideae and B. phytophthorus, 834–835; western yellow blight, or yellows, 709–710; yellows or tomato curly top, 709–710
Tomatoes, "streak" of, produced by disturbing principle from healthy potatoes in combination with tomato-mosaic virus, 311
TOMPKINS, C. M., and S. B. NUCKOLS, Development of storage diseases in sugar beets resulting from hook injury, 939–941
Topography in relation to severity of stem-rust epidemics, 90–91
Toxicity, lime-sulphur, nature, 355–360
Transmission, mechanical, of mosaic of Lilium auratum, 1025–1026; of bean mosaic, 155; of black leg by seed-corn maggot, 459; of cucumber mosaic by aphis, 143; of potato spindle tuber by grasshoppers, 445–448; of potato spindle tuber by grasshoppers (Locustidae), 140; of the virus of curly top sugar beets through different solutions, 675–679; soil, of tomato mosaic and streak, 155
Transplants, studies on crown gall, 130
Tranzschelia punctata, action of lime sulphur on spores, 355–360
Treatment, of millet seed to prevent smut, 479; of soils infested with Plasmodiophora brassicae, 141–142; of sweet potatoes for black-rot control, 153–154
Treatments, dust, for oat smuts, 139
Trees, nursery, apple and quince, overgrowths and hairy root on, 126
Trichoderma lignorum, 949
Trifolium, hybridum, Uromyces trifolii on,

experimental control of teliospore and urediniospore formation, 172–175, 188–191, 206; pratense, lesions produced by Pseudopeziza trifolii, 742

Triticum, vulgare, inheritance of resistance to Puccinia graminis in a cross between two varieties of, 631–658; vulgare, P. triticina on, experimental control of teliospore and urediniospore formation, 179, 195–196

Turf diseases, control of, with mercury, 137

Turnip, susceptibility to Bacillus carotovorus, B. aroideae, and B. phytophthorus, 833–835

Tussilago farfara, lesions produced by Sphaerella tussilaginis, 745–746

Uniform wheat-rust nurseries, 13–17, 20–31

Urediniospore formation, experimental control of, 157–213

Uredinopsis, copelandi, on Athyrium cyclosorum and Felix bulbifera, 705–708; macrosperma, on Pteridium aquilinum and P. pubescens, 705–708; mirabilis, on Woodwardia areolata and Onoclea sensibilis, 705–708; osmundae, on Osmunda species, 705–708; phegopteridis, on Phegopteris dryopteris, 705–708; struthiopteridis, on Woodwardia virginica and Struthiopteris germanica, 705–708

Uredo alpestris, 670

Uromyces, appendiculatus on Phaseolus vulgaris, experimental control of teliospore and urediniospore formation, 163–169, 183, 184–187, 206; polygoni on Polygonum aviculare, experimental control of teliospore and urediniospore formation, 178–179, 206; scillarum, 672; trifolii on Trifolium hybridum, experimental control of teliospore and urediniospore formation, 172–175, 188–191, 206

Ustilago, avenae, cultural differences between physiologic forms of, 969–972; avenae, effect of temperature on growth of physiologic forms of, 981; avenae, varietal resistance of oats to, 993–997; crameri, 911–912; hordei, cultural differences between physiologic forms of, 964–968; hordei, effect of temperature on growth of physiologic forms of, 979–981; hordei, pathogenicity of physiologic forms of, 992–993; levis, cultural differences between physiologic forms of, 969; levis, effect of temperature on growth of physiologic forms of, 981; levis, varietal resistance of oats to, 993–997; nuda, cultural differences between physiologic forms of, 962–964; nuda, effect of temperature on growth of physiologic forms of, 976–979; striaeformis, two physiologic forms of, 149; tritici, cultural differences between physiologic forms of, 960–962; tritici, effect of temperature on growth of physiologic forms of, 973–976; zeae, influence of environmental factors on seasonal prevalence, 589–598; zeae, losses due to smut infection in selfed lines of corn, 599–602

Vaccinium, mycorrhizal fungus, of, 148; species, alternate hosts of fir-needle rusts, 705–708

VALLEAU, W. D., and E. M. JOHNSON, Experiments and observations on the control of true tobacco mosaic, 132; Some virus diseases of tobacco in Kentucky, 132–133

van der BIJL, P. A., see BIJL, P. A. van der

Varietal, resistance of watermelons to wilt (Fusarium niveum), 144; susceptibility of potatoes to Fusarium wilt and stem-end rot, 307–309

Velvet bean, bacterial spot of, 460

Ventilation, forced, as a means of controlling tomato Cladosporium and Septoria in hotbeds, 1027–1029

Venturia, cerasi, 328; inaequalis, 943–944; inaequalis, certain sulphur fungicides in the control of apple scab, 146–147; inaequalis, effect of environment upon, 375–418; inaequalis, factors important in development of perithecia of, 145–146; inaequalis, fall application of fungicides in relation to apple-scab control, 146

Verbascum blattaria, lesions produced by Septoria verbasicola, 733–734

Verticillium, albo-atrum, reaction of plant

stems to products of, 604–605; wilt of heliotrope, 246
Vinca rosea, occurrence of intracellular bodies in spike disease of, 815–817
Virus, beet curly-top, transmission through different solutions, 675–679; disease, of tobacco, ring spot, 133; diseases observed by the Allison V. Armour Expedition, 155; diseases, of tobacco in Kentucky, 132–133; of sugar beet curly top, 951; tobacco-mosaic, accuracy in comparing various concentrations of, 132; tomato-mosaic, a streak of tomatoes produced by disturbing principle from healthy potatoes in combination with, 311
Viruses, plant, studies on attenuation of, 156

WALKER, J. C., and F. L. WELLMAN, A Fusarium-resistant cabbage of Jersey Wakefield type, 142
WALKER, M. N., see DOOLITTLE, S. P.
Walnut, crown rot, of, 948–949
Warty disease, or black scab, of potato, in Belgium, 126
Washing seed, effect on wheat-stinking-smut infection, 711–713
Washingtonia robusta, leaf spot due to Cylindrocladium macrosporum, 219–225
Watermelon, studies with wilt of, caused by Fusarium niveum, 143–144; varieties of, resistant to wilt (Fusarium niveum), 144
WATERS, CHARLES W., The control of teliospore and urediniospore formation by experimental methods, 157–213
Weather in relation to severity of stem-rust epidemics, 91–105
WEBER, GEORGE F., Biography of OWEN, F. BURGER, 627–630; Cucumber fruit rot and angular leaf spot, 133
WELLMAN, F. L., The reaction and treatment of soils infested with Plasmodiophora brassicae Wor., 141–142; see WALKER, J. C.
WELSH, J. N., see GOULDEN, C. H.
Western yellow blight of tomatoes, or yellows, 709–710
Wheat, effect of ecological factors on susceptibility of varieties of, to Puccinia graminis tritici, 81–108; effect of washing seed on stinking-smut infection, 711–713; inheritance of resistance to Puccinia graminis tritici in a cross between two varieties of, 631–658; new physiologic forms of Tilletia tritici in, 139; physiologic forms of stem rust in Canada, 478–479; Puccinia triticina on, experimental control of teliospore and urediniospore formation, 179, 195–196; reaction to take-all in Canada, 477; rust epidemics, 369–374; smut, physiologic forms of, 579–588; spring, varietal resistance to Tilletia levis, 981–984
WHETZEL, H. H., The crown-elongation disease of the peony, 243–244
White spot of alfalfa, caused by irrigation, 136–137
WILLIAMS, ROBERT, see YOUNG, H. C.
WILSON, E. E., Factors important in the development of perithecia of Venturia inaequalis, 145–146; see KEITT, G. W.; Studies of the ascigerous stage of Venturia inaequalis (Cke.) Wint. in relation to certain factors of the environment, 375–418
Wilt, Fusarium, and stem-end rot, varietal susceptibility of potatoes to, 307–309; (Fusarium niveum), of watermelon, varietal resistance to, 144; Fusarium, of China aster, 150; relation of fungous products to, 603–610; Verticillium of heliotrope, 246; watermelon, studies of, 143–144
Wilt-resistant cotton varieties, 134
WINGARD, S. A., and F. D. FROMME, Tobacco ring spot: A virus disease with a wide host range, 133
WOLF, FREDERICK A., Further observations on Corticium koleroga (Cke.) v. Hohn., 147–148
WOODROOF, NAOMI CHAPMAN, Cotton-seed treatment by the dusting method, 134
Woodwardia, areolata, host of Uredinopsis mirabilis, 705–708; virginica, host of U. struthiopteridis, 705–708
Wound overgrowth, correlation of, and crown gall of apple, 128
Wound overgrowths, studies of the history of development of, on apple grafts and

of the influence of wrappers on their suppression, 128

Wrappers, influence of, on the suppression of wound overgrowths, on apple grafts, 128

WRIGHT, WILLIAM H., and SKORIC, VLADIMIR, Demonstration of bacteria in plant tissues by means of Giemsa stain, 803–807

Xylaria mali, cause of black root rot of apple, 145

Yellows, aster, studies on host range of, 156; of potato, a new and destructive disease in Utah, 140–141; tomato, 709–710, 950

YOUNG, H. C., and ROBERT WILLIAMS, Factors affecting the fungicidal property of sulphur, 147

YOUNG, V. H., Notice of field meeting of the Southern Section of the American Phytopathological Society, 460–461

YU, T. F., see PORTER, R. H.

Zantedeschia aethiopica, susceptibility to Bacillus aroideae, 835–836

Zea mays, host of Pythium arrhenomanes, 873–875; Puccinia sorghi on, experimental control of teliospore and uredinio-spore formation, 179–180, 194–195, 206; see corn